

PATENTS
514453-3916


AMENDMENT

IN THE CLAIMS

Please enter the following amendments to the claims without prejudice, without admission, without surrender of subject matter, and without any intention of creating any estoppel as to equivalents as follows:

Claim 1 (currently amended)

1. A five-membered ring compound of the formula (I),



where the symbols and indices have the following meanings:

E is a radical T-Z-R² containing a five-membered ring, where:

- (i) T is undirected and is

4-fluorothiophene-2,5-diyl, 3-fluorothiophene-2,5-diyl,
3-fluorothiophene-2,4-diyl or 5-fluorothiophene-2,4-diyl

Z is a single bond or -O-

R² is hydrogen or a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one nonterminal CH₂ group may be replaced by -O- or -OC(=O)- or -C(=O)O- and/or one or more H atoms may be replaced by F, with the provisos that
a) the -CH₂- group nearest to the thiophene cannot be replaced by -O- when Z is -O-
b) R² can only be hydrogen when Z is a single bond,

Y is -OC(=O)-, -OCH₂-, -CH₂CH₂-

a, b are each, independently of one another, 0 or 1

- (ii) T is furan-2,5-diyl or furan-2,4-diyl

Z is a single bond or -O-

R² is a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one nonterminal CH₂

PATENTS
514453-3916

group nonadjacent to furan may be replaced by -O- or -OC(=O)- or
-C(=O)O- and/or one or more H atoms may be replaced by F,

Y is -OC(=O)-, -OCH₂-, -CH₂CH₂-

a, b are each, independently of one another, 0 or 1

- (iii) **T** is undirected and is isoxazole-3,5-diyl
Z is a single bond or -O-
R² is hydrogen or a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one nonterminal CH₂ group may be replaced by -O- or -OC(=O)- or -C(=O)O- and/or one or more H atoms may be replaced by F, with the provisos that
a) the -CH₂- group nearest to the isoxazole cannot be replaced by -O- when Z is -O-
b) **R²** can only be hydrogen when Z is a single bond,

a is 1.

b is 0 or 1

Y is -OC(=O)-, -OCH₂-, -CH₂CH₂-

- (iv) **T** is undirected and is thiazole-2,5-diyl or thiazole-2,4-diyl
Z is a single bond
R² is hydrogen or a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one nonterminal CH₂ group may be replaced by -O- or -OC(=O)- or -C(=O)O- and/or one or more H atoms may be replaced by F,


Y is -OC(=O)-, -OCH₂-, -CH₂CH₂-

a, b are each, independently of one another, 0 or 1

- (v) **T** is cyclopentane-1,3-diyl
Z is a single bond or -O-

PATENTS
514453-3916

~~R² is hydrogen or a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one nonterminal CH₂ group may be replaced by -O- or -OC(=O)- or -C(=O)O- and/or one or more H atoms may be replaced by F, with the provisos that~~
~~a) the -CH₂- group nearest to the cyclopentane cannot be replaced by -O- when Z is -O-~~
~~b) R² can only be hydrogen when Z is a single bond,~~
 with the exception of compounds of the formula (II)



in which

~~R⁴ is as defined for R²~~

~~G is trans-4-propyl-cyclohexyl or trans-4-butyl-cyclohexyl or an alkyl group of 1 to 15 carbon atoms, in which, in addition, one or more nonadjacent CH₂ groups may be replaced by -O-, -CO-, -OCO-, -O-CO-O-, -CHhalogen-, -CHCN- and/or -CH=CH- or is F, CN,~~

(v)(vi) T is cyclopentane-1,3-diyl, in which one -CH₂CH₂- or -CH₂CH- group is replaced by a -CH=CH- or -CH=C- group respectively

Z is a single bond

R² is hydrogen or a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one nonterminal CH₂ group may be replaced by -O- or -OC(=O)- or -C(=O)O- and/or one or more H atoms may be replaced by F, with the proviso that the -CH₂- group nearest to the cyclopentene cannot be replaced and where

Y cannot be -CH₂-CH₂-,

PATENTS
514453-3916

a is 1

b is 0 or 1

Y is -OC(=O)-, -OCH₂-

R¹ is hydrogen or a straight-chain or branched C₁₋₂₀-alkyl or C₂₋₂₀-alkenyl radical (with or without asymmetric carbon atoms), where

- a) one or two nonterminal CH₂ groups may be replaced, independently of one another, by -O- or -C(=O)-, with the proviso that two adjacent CH₂ groups cannot be replaced in the same way, and/or
- b) one CH₂ group may be replaced by -C≡C-, and/or
- c) one CH₂ group may be replaced by -Si(CH₃)₂-, cyclopropane-1,2-diyl, cyclobutane-1,3-diyl, cyclopentane-1,4-diyl, bicyclo[1.1.1]pentane-1,3-diyl or cyclohexane-1,4-diyl, and/or
- d) one or more H atoms may be replaced by F and/or CN,
- e) in the case of a branched alkyl radical containing asymmetric carbon atoms, the asymmetric carbon atoms have -CH₃, -OCH₃, -CF₃, F, CN and/or Cl as substituents or are incorporated into a 3- to 7-membered ring, in which, in addition, one or two non-adjacent CH₂ groups may be replaced by -O- and one CH₂ group non-adjacent to these groups may be replaced by -OC(=O)-;

X is a single bond, -O-, OC(=O)-, -C(=O)O- or -OC(=O)O-

Y is ~~OC(-O)-, OCH₂-, CH₂CH₂-~~

A¹, A², A³ are each, independently of one another, phenylene-1,4-diyl, unsubstituted or monosubstituted or disubstituted by CN or F, phenylene-1,3-diyl, unsubstituted or monosubstituted or disubstituted by CN or F, cyclohexane-1,4-diyl, in which one or two H atoms may be replaced by CN and/or CH₃ and/or F, 1-cyclohexene-1,4-diyl, in which one H atom may be replaced by F, 1-alkyl-1-silacyclohexane-1,4-diyl, pyridine-2,5-diyl, unsubstituted or monosubstituted by F, pyrimidine-2,5-

BEST AVAILABLE COPY

PATENTS
514453-3916

diyl, unsubstituted or monosubstituted by F, cyclopentane-2,5-diyl or thiophene-2,5-diyl;

M^1 , M^2 are undirected and are each, independently of one another, $-\text{OC}(=\text{O})-$,

$-\text{OCH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{OC}(=\text{O})\text{CH}_2\text{CH}_2-$, $-\text{OCH}_2\text{CH}_2\text{CH}_2-$, $-\text{C}\equiv\text{C}-$,

$-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$ or a single bond;

~~a , b are each, independently of one another, 0 or 1.~~

Claim 2 (previously presented)

2. A liquid-crystal mixture comprising at least one compound of the formula (I) as claimed in claim 1.

Claim 3 (currently amended)

3. A liquid-crystal mixture as claimed in claim 2, which comprises from 0.01 to 80% by weight, based on the entire weight of the mixture, of one or more compounds of the formula (I).

Claim 4 (previously presented)

4. A liquid-crystal mixture as claimed in claim 2, which is ferroelectric (chiral smectic).

Claim 5 (previously presented)

5. A liquid-crystal mixture as claimed in claim 2, which is nematic.

Claim 6 (previously presented)

6. A ferroelectric switching and/or display device, which contains a ferroelectric liquid-crystal mixture as claimed in claim 4.

Claim 7 (previously presented)

7. A ferroelectric switching and/or display device as claimed in claim 6, which contains active matrix elements and wherein the liquid-crystal layer forms a monostable monodomain.

BEST AVAILABLE COPYPATENTS
514453-3916

Claims 8-12 (cancelled)

Claim 13 (new)

13. A liquid-crystal mixture as claimed in claim 2, which comprises 0.1 to 30% by weight, based on the entire weight of the mixture, of one or more compounds of formula (I).

Claim 14 (new)


14. The liquid-crystal mixture of claim 13, which is ferroelectric (chiral smectic) and further comprises one or more compounds having a smectic phase.

Claim 15 (new)

15. The liquid-crystal mixture of claim 13, which is nematic and further comprises one or more compounds having a nematic phase.

Claim 16 (new)

16. The ferroelectric switching and/or display device of claim 7, which contains a liquid-crystal mixture comprising at least one compound of the formula (I):



where the symbols and indices have the following meanings:

E is a radical T-Z-R² containing a five-membered ring, where:

- (i) T is undirected and is

4-fluorothiophene-2,5-diyl, 3-fluorothiophene-2,5-diyl,
3-fluorothiophene-2,4-diyl or 5-fluorothiophene-2,4-diyl

Z is a single bond or -O-

R² is hydrogen or a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one nonterminal CH₂ group may be replaced by -O- or -OC(=O)- or -C(=O)O- and/or one or more H atoms may be replaced by F, with the provisos that
a) the -CH₂- group nearest to the thiophene cannot be replaced by -O- when Z is -O-

BEST AVAILABLE COPY

PATENTS
514453-3916b) R^2 can only be hydrogen when Z is a single bond,Y is $-OC(=O)-$, $-OCH_2-$, $-CH_2CH_2-$

a, b are each, independently of one another, 0 or 1

(ii) T is furan-2,5-diyl or furan-2,4-diyl

Z is a single bond or $-O-$ R^2 is a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one nonterminal CH_2 group nonadjacent to furan may be replaced by $-O-$ or $-OC(=O)-$ or $-C(=O)O-$ and/or one or more H atoms may be replaced by F,Y is $-OC(=O)-$, $-OCH_2-$, $-CH_2CH_2-$

a, b are each, independently of one another, 0 or 1

(iii) T is undirected and is isoxazole-3,5-diyl

Z is a single bond or $-O-$ R^2 is hydrogen or a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one nonterminal CH_2 group may be replaced by $-O-$ or $-OC(=O)-$ or $-C(=O)O-$ and/or one or more H atoms may be replaced by F, with the provisos that
a) the $-CH_2-$ group nearest to the isoxazole cannot be replaced by $-O-$ when Z is $-O-$ b) R^2 can only be hydrogen when Z is a single bond,

a is 1

b is 0 or 1

Y is $-OC(=O)-$, $-OCH_2-$, $-CH_2CH_2-$

(iv) T is undirected and is thiazole-2,5-diyl or thiazole-2,4-diyl

Z is a single bond

BEST AVAILABLE COPY

PATENTS
514453-3916

R² is hydrogen or a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one nonterminal CH₂ group may be replaced by -O- or -OC(=O)- or -C(=O)O- and/or one or more H atoms may be replaced by F,

Y is -OC(=O)-, -OCH₂-, -CH₂CH₂-

a, b are each, independently of one another, 0 or 1

(v) **T** is cyclopentane-1,3-diyl, in which one -CH₂CH₂- or -CH₂CH- group is replaced by a -CH=CH- or CH=C- group respectively

Z is a single bond

R² is hydrogen or a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one nonterminal CH₂ group may be replaced by -O- or -OC(=O)- or -C(=O)O- and/or one or more H atoms may be replaced by F, with the proviso that the -CH₂- group nearest to the cyclopentene cannot be replaced and where

Y cannot be -CH₂-CH₂-,

a is 1

b is 0 or 1

Y is -OC(=O)-, -OCH₂-

R¹ is hydrogen or a straight-chain or branched C₁₋₂₀-alkyl or C₂₋₂₀-alkenyl radical (with or without asymmetric carbon atoms), where

a) one or two nonterminal CH₂ groups may be replaced, independently of one another, by -O- or -C(=O)-, with the proviso that two adjacent CH₂ groups cannot be replaced in the same way, and/or

b) one CH₂ group may be replaced by -C≡C-, and/or

PATENTS
514453-3916

- c) one CH₂ group may be replaced by -Si(CH₃)₂-, cyclopropane-1,2-diyl, cyclobutane-1,3-diyl, cyclopentane-1,4-diyl, bicyclo[1.1.1]pentane-1,3-diyl or cyclohexane-1,4-diyl, and/or
- d) one or more H atoms may be replaced by F and/or CN,
- e) in the case of a branched alkyl radical containing asymmetric carbon atoms, the asymmetric carbon atoms have -CH₃, -OCH₃, -CF₃, F, CN and/or Cl as substituents or are incorporated into a 3- to 7-membered ring, in which, in addition, one or two non-adjacent CH₂ groups may be replaced by -O- and one CH₂ group non-adjacent to these groups may be replaced by -OC(=O)-;

X is a single bond, -O-, OC(=O)-, -C(=O)O- or -OC(=O)O-


A¹, A², A³ are each, independently of one another, phenylene-1,4-diyl, unsubstituted or monosubstituted or disubstituted by CN or F, phenylene-1,3-diyl, unsubstituted or monosubstituted or disubstituted by CN or F, cyclohexane-1,4-diyl, in which one or two H atoms may be replaced by CN and/or CH₃ and/or F, 1-cyclohexene-1,4-diyl, in which one H atom may be replaced by F, 1-alkyl-1-silacyclohexane-1,4-diyl, pyridine-2,5-diyl, unsubstituted or monosubstituted by F, pyrimidine-2,5-diyl, unsubstituted or monosubstituted by F, cyclopentane-2,5-diyl or thiophene-2,5-diyl;

M¹, M² are undirected and are each, independently of one another, -OC(=O)-, -OCH₂-, -CH₂CH₂-, -OC(=O)CH₂CH₂-, -OCH₂CH₂CH₂-, -C≡C-, -CH₂CH₂CH₂CH₂- or a single bond;

wherein said liquid crystal mixture is ferroelectric (chiral smectic) and further comprises one or more compounds having a smectic phase.

Claim 17 (new)

17. The ferroelectric switching and/or display device of claim 7, which contains a liquid-crystal mixture comprising at least one compound of the formula (I):



PATENTS
514453-3916

where the symbols and indices have the following meanings:

E is a radical $T-Z-R^2$ containing a five-membered ring, where:

(i) **T** is undirected and is

4-fluorothiophene-2,5-diyl, 3-fluorothiophene-2,5-diyl,
3-fluorothiophene-2,4-diyl or 5-fluorothiophene-2,4-diyl

Z is a single bond or -O-

R² is hydrogen or a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one nonterminal CH_2 group may be replaced by -O- or -OC(=O)- or -C(=O)O- and/or one or more H atoms may be replaced by F, with the provisos that
a) the - CH_2 - group nearest to the thiophene cannot be replaced by -O- when Z is -O-
b) R^2 can only be hydrogen when Z is a single bond,

Y is -OC(=O)-, -OCH₂-, -CH₂CH₂-

a, b are each, independently of one another, 0 or 1

(ii) **T** is furan-2,5-diyl or furan-2,4-diyl

Z is a single bond or -O-

R² is a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one nonterminal CH_2 group nonadjacent to furan may be replaced by -O- or -OC(=O)- or -C(=O)O- and/or one or more H atoms may be replaced by F,

Y is -OC(=O)-, -OCH₂-, -CH₂CH₂-

a, b are each, independently of one another, 0 or 1

(iii) **T** is undirected and is isoxazole-3,5-diyl

Z is a single bond or -O-

R² is hydrogen or a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one

PATENTS
514453-3916

nonterminal CH₂ group may be replaced by -O- or -OC(=O)- or -C(=O)O- and/or one or more H atoms may be replaced by F, with the provisos that
a) the -CH₂- group nearest to the isoxazole cannot be replaced by -O- when Z is -O-

b) R² can only be hydrogen when Z is a single bond,

a is 1

b is 0 or 1

Y is -OC(=O)-, -OCH₂-, -CH₂CH₂-

(iv) T is undirected and is thiazole-2,5-diyl or thiazole-2,4-diyl

Z is a single bond

R² is hydrogen or a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one nonterminal CH₂ group may be replaced by -O- or -OC(=O)- or -C(=O)O- and/or one or more H atoms may be replaced by F,

Y is -OC(=O)-, -OCH₂-, -CH₂CH₂-

a, b are each, independently of one another, 0 or 1

(v) T is cyclopentane-1,3-diyl, in which one -CH₂CH₂- or -CH₂CH- group is replaced by a -CH=CH- or -CH=C- group respectively

Z is a single bond

R² is hydrogen or a straight-chain or branched alkyl radical (with or without asymmetric carbon atoms) having 1 to 20 carbon atoms, where one nonterminal CH₂ group may be replaced by -O- or -OC(=O)- or -C(=O)O- and/or one or more H atoms may be replaced by F, with the proviso that the -CH₂- group nearest to the cyclopentene cannot be replaced and where

Y cannot be -CH₂-CH₂-,

a is 1

PATENTS
514453-3916**b** is 0 or 1**Y** is -OC(=O)-, -OCH₂-**R¹** is hydrogen or a straight-chain or branched C₁₋₂₀-alkyl or C₂₋₂₀-alkenyl radical (with or without asymmetric carbon atoms), where

- a) one or two nonterminal CH₂ groups may be replaced, independently of one another, by -O- or -C(=O)-, with the proviso that two adjacent CH₂ groups cannot be replaced in the same way, and/or
- b) one CH₂ group may be replaced by -C≡C-, and/or
- c) one CH₂ group may be replaced by -Si(CH₃)₂-, cyclopropane-1,2-diyl, cyclobutane-1,3-diyl, cyclopentane-1,4-diyl, bicyclo[1.1.1]pentane-1,3-diyl or cyclohexane-1,4-diyl, and/or
- d) one or more H atoms may be replaced by F and/or CN,
- e) in the case of a branched alkyl radical containing asymmetric carbon atoms, the asymmetric carbon atoms have -CH₃, -OCH₃, -CF₃, F, CN and/or Cl as substituents or are incorporated into a 3- to 7-membered ring, in which, in addition, one or two non-adjacent CH₂ groups may be replaced by -O- and one CH₂ group non-adjacent to these groups may be replaced by -OC(=O)-;

X is a single bond, -O-, OC(=O)-, -C(=O)O- or -OC(=O)O-

A¹, A², A³ are each, independently of one another, phenylene-1,4-diyl, unsubstituted or monosubstituted or disubstituted by CN or F, phenylene-1,3-diyl, unsubstituted or monosubstituted or disubstituted by CN or F, cyclohexane-1,4-diyl, in which one or two H atoms may be replaced by CN and/or CH₃ and/or F, 1-cyclohexene-1,4-diyl, in which one H atom may be replaced by F, 1-alkyl-1-silacyclohexane-1,4-diyl, pyridine-2,5-diyl, unsubstituted or monosubstituted by F, pyrimidine-2,5-diyl, unsubstituted or monosubstituted by F, cyclopentane-2,5-diyl or thiophene-2,5-diyl;

PATENTS
514453-3916

M^1, M^2 are undirected and are each, independently of one another, $-OC(=O)-$,
 $-OCH_2-$, $-CH_2CH_2-$, $-OC(=O)CH_2CH_2-$, $-OCH_2CH_2CH_2-$, $-C\equiv C-$,
 $-CH_2CH_2CH_2CH_2-$ or a single bond;

wherein said liquid crystal mixture is nematic and further comprises one or more
compounds having a nematic phase.